



Left: Highvale visitant. (R): Flowers (top); aerial tubers (bottom).

<i>Common Name</i>	<i>Botanical Name</i>	<i>Treatment</i>
Madeira vine	<i>Anredera cordifolia</i>	<ul style="list-style-type: none"> <li>• Dig small tuberlings and tubers, bag and dispose in rubbish - not garden waste - or spray plants in place with <b>1:50 Starane</b>.</li> <li>• Cut stem and dip both ends into <b>1:5 glyphosate or 1:5 Starane</b> in bottle. Must be left for a few weeks to destroy tubers on vine. This must be placed well out of reach of children and animals.</li> </ul>

### Notes:

- The most effective time to apply herbicide is during times of vigorous plant growth. It is important to kill weeds before they produce viable reproductive entities such as seed, bulbs, etc.
- Wetting agents such as Chemwet or Protec can help the efficiency of the sprayed herbicide.
- After initial weed clearing when native regrowth has begun to reappear it is best to hand pull or dig out new weeds.
- Utmost care must be taken to protect children, animals and desirable plants from herbicides, including from overspray.
- Bagged weeds must be disposed of in general rubbish, which will go to landfill. Do **NOT** put weeds in garden waste.
- Many garden plants have become weeds by escaping backyards. To avoid adding to this you need to firstly recognise which plants have become weeds, then look for a suitable, non-invasive alternative plant. The safest is usually a native Australian species.